

Search for the rare decay $\eta' \rightarrow \pi^0 \pi^0 \pi^0 \pi^0$ at BESIII

M. Ablikim,¹ M. N. Achasov,^{10,†} P. Adlarson,⁶³ S. Ahmed,¹⁵ M. Albrecht,⁴ M. Alekseev,^{62a,62c} A. Amoroso,^{62a,62c} Q. An,^{59,47} Y. Bai Anita,^{21,46} O. Bakina,²⁸ R. Baldini Ferroli,^{23a} I. Balossino,^{24a} Y. Ban,^{37,‡} K. Begzsuren,²⁶ J. V. Bennett,⁵ N. Berger,²⁷ M. Bertani,^{23a} D. Bettoni,^{24a} F. Bianchi,^{62a,62c} J. Biernat,⁶³ J. Bloms,⁵⁶ I. Boyko,²⁸ R. A. Briere,⁵ H. Cai,⁶⁴ X. Cai,^{1,47} A. Calcaterra,^{23a} G. F. Cao,^{1,51} N. Cao,^{1,51} S. A. Cetin,^{50b} J. Chai,^{62c} J. F. Chang,^{1,47} W. L. Chang,^{1,51*} G. Chelkov,^{28,§,||} D. Y. Chen,⁶ G. Chen,¹ H. S. Chen,^{1,51} J. Chen,¹⁶ M. L. Chen,^{1,47} S. J. Chen,³⁵ X. R. Chen,²⁵ Y. B. Chen,^{1,47} W. Cheng,^{62c} G. Cibinetto,^{24a} F. Cossio,^{62c} X. F. Cui,³⁶ H. L. Dai,^{1,47} J. P. Dai,^{41,¶} X. C. Dai,^{1,51} A. Dbeysy,¹⁵ D. Dedovich,²⁸ Z. Y. Deng,¹ A. Denig,²⁷ I. Denysenko,²⁸ M. Destefanis,^{62a,62c} F. De Mori,^{62a,62c} Y. Ding,³³ C. Dong,³⁶ J. Dong,^{1,47} L. Y. Dong,^{1,51} M. Y. Dong,^{1,47,51} Z. L. Dou,³⁵ S. X. Du,⁶⁷ J. Fang,^{1,47} S. S. Fang,^{1,51} Y. Fang,¹ R. Farinelli,^{24a,24b} L. Fava,^{62b,62c} F. Feldbauer,⁴ G. Felici,^{23a} C. Q. Feng,^{59,47} M. Fritsch,⁴ C. D. Fu,¹ Y. Fu,¹ X. L. Gao,^{59,47} Y. Gao,^{37,‡} Y. Gao,⁶⁰ Y. G. Gao,⁶ I. Garzia,^{24a,24b} E. M. Gersabeck,⁵⁴ A. Gilman,⁵⁵ K. Goetzen,¹¹ L. Gong,³⁶ W. X. Gong,^{1,47} W. Gradl,²⁷ M. Greco,^{62a,62c} L. M. Gu,³⁵ M. H. Gu,^{1,47} S. Gu,² Y. T. Gu,¹³ A. Q. Guo,²² L. B. Guo,³⁴ R. P. Guo,³⁹ Y. P. Guo,²⁷ A. Guskov,²⁸ S. Han,⁶⁴ T. T. Han,⁴⁰ X. Q. Hao,¹⁶ F. A. Harris,⁵² K. L. He,^{1,51} F. H. Heinsius,⁴ T. Held,⁴ Y. K. Heng,^{1,47,51} M. Himmelreich,^{11,**} T. Holtmann,⁴ Y. R. Hou,⁵¹ Z. L. Hou,¹ H. M. Hu,^{1,51} J. F. Hu,^{41,¶} T. Hu,^{1,47,51} Y. Hu,¹ G. S. Huang,^{59,47} J. S. Huang,¹⁶ X. T. Huang,⁴⁰ X. Z. Huang,³⁵ N. Huesken,⁵⁶ T. Hussain,⁶¹ W. Ikegami Andersson,⁶³ W. Imoehl,²² M. Irshad,^{59,47} S. Jaeger,⁴ Q. Ji,¹ Q. P. Ji,¹⁶ X. B. Ji,^{1,51} X. L. Ji,^{1,47} H. B. Jiang,⁴⁰ X. S. Jiang,^{1,47,51} X. Y. Jiang,³⁶ J. B. Jiao,⁴⁰ Z. Jiao,¹⁸ D. P. Jin,^{1,47,51} S. Jin,³⁵ Y. Jin,⁵³ T. Johansson,⁶³ N. Kalantar-Nayestanaki,³⁰ X. S. Kang,³³ R. Kappert,³⁰ M. Kavatsyuk,³⁰ B. C. Ke,^{42,1} I. K. Keshk,⁴ A. Khoukaz,⁵⁶ P. Kiese,²⁷ R. Kiuchi,¹ R. Kliemt,¹¹ L. Koch,²⁹ O. B. Kolcu,^{50b,††} B. Kopf,⁴ M. Kuemmel,⁴ M. Kuessner,⁴ A. Kupsc,⁶³ M. G. Kurth,^{1,51} W. Kühn,²⁹ J. S. Lange,²⁹ P. Larin,¹⁵ L. Lavezzi,^{62c} H. Leithoff,²⁷ T. Lenz,²⁷ C. Li,³⁸ C. H. Li,³² Cheng Li,^{59,47} D. M. Li,⁶⁷ F. Li,^{1,47} G. Li,¹ H. B. Li,^{1,51} H. J. Li,^{9,‡‡} J. C. Li,¹ Ke Li,¹ L. K. Li,¹ Lei Li,³ P. L. Li,^{59,47} P. R. Li,³¹ W. D. Li,^{1,51} W. G. Li,¹ X. H. Li,^{59,47} X. L. Li,⁴⁰ X. N. Li,^{1,47} Z. B. Li,⁴⁸ Z. Y. Li,⁴⁸ H. Liang,^{1,51} H. Liang,^{59,47} Y. F. Liang,⁴⁴ Y. T. Liang,²⁵ G. R. Liao,¹² L. Z. Liao,^{1,51} J. Libby,²¹ C. X. Lin,⁴⁸ D. X. Lin,¹⁵ Y. J. Lin,¹³ B. Liu,^{41,¶} B. J. Liu,¹ C. X. Liu,¹ D. Liu,^{59,47} D. Y. Liu,^{41,¶} F. H. Liu,⁴³ Fang Liu,¹ Feng Liu,⁶ H. B. Liu,¹³ H. M. Liu,^{1,51} Huanhuan Liu,¹ Huihui Liu,¹⁷ J. B. Liu,^{59,47} J. Y. Liu,^{1,51} K. Liu,¹ K. Y. Liu,³³ Ke Liu,⁶ L. Liu,^{59,47} L. Y. Liu,¹³ Q. Liu,⁵¹ S. B. Liu,^{59,47} T. Liu,^{1,51} X. Liu,³¹ X. Y. Liu,^{1,51} Y. B. Liu,³⁶ Z. A. Liu,^{1,47,51} Z. Q. Liu,⁴⁰ Y. F. Long,^{37,‡} X. C. Lou,^{1,47,51} H. J. Lu,¹⁸ J. D. Lu,^{1,51} J. G. Lu,^{1,47} Y. Lu,¹ Y. P. Lu,^{1,47} C. L. Luo,³⁴ M. X. Luo,⁶⁶ P. W. Luo,⁴⁸ T. Luo,^{9,‡‡} X. L. Luo,^{1,47} S. Lusso,^{62c} X. R. Lyu,⁵¹ F. C. Ma,³³ H. L. Ma,¹ L. L. Ma,⁴⁰ M. M. Ma,^{1,51} Q. M. Ma,¹ R. T. Ma,⁵¹ X. N. Ma,³⁶ X. X. Ma,^{1,51} X. Y. Ma,^{1,47} Y. M. Ma,⁴⁰ F. E. Maas,¹⁵ M. Maggiora,^{62a,62c} S. Maldaner,²⁷ S. Malde,⁵⁷ Q. A. Malik,⁶¹ A. Mangoni,^{23b} Y. J. Mao,^{37,‡} Z. P. Mao,¹ S. Marcello,^{62a,62c} Z. X. Meng,⁵³ J. G. Messchendorp,³⁰ G. Mezzadri,^{24a} J. Min,^{1,47} T. J. Min,³⁵ R. E. Mitchell,²² X. H. Mo,^{1,47,51} Y. J. Mo,⁶ C. Morales Morales,¹⁵ N. Yu. Muchnoi,^{10,†} H. Muramatsu,⁵⁵ A. Mustafa,⁴ S. Nakhoul,^{11,**} Y. Nefedov,²⁸ F. Nerling,^{11,**} I. B. Nikolaev,^{10,†} Z. Ning,^{1,47} S. Nisar,^{8,§§} S. L. Olsen,⁵¹ Q. Ouyang,^{1,47,51} S. Pacetti,^{23b} Y. Pan,^{59,47} M. Papenbrock,⁶³ P. Patteri,^{23a} M. Pelizaeus,⁴ H. P. Peng,^{59,47} K. Peters,^{11,**} J. Pettersson,⁶³ J. L. Ping,³⁴ R. G. Ping,^{1,51} A. Pitka,⁴ R. Poling,⁵⁵ V. Prasad,^{59,47} H. Qi,^{59,47} M. Qi,³⁵ T. Y. Qi,² S. Qian,^{1,47} C. F. Qiao,⁵¹ X. P. Qin,¹³ X. S. Qin,⁴ Z. H. Qin,^{1,47} J. F. Qiu,¹ S. Q. Qu,³⁶ K. H. Rashid,⁶¹ K. Ravindran,²¹ C. F. Redmer,²⁷ M. Richter,⁴ A. Rivetti,^{62c} V. Rodin,³⁰ M. Rolo,^{62c} G. Rong,^{1,51} Ch. Rosner,¹⁵ M. Rump,⁵⁶ A. Sarantsev,^{28,|||} M. Savrié,^{24b} Y. Schelhaas,²⁷ C. Schnier,⁴ K. Schoenning,⁶³ W. Shan,¹⁹ X. Y. Shan,^{59,47} M. Shao,^{59,47} C. P. Shen,² P. X. Shen,³⁶ X. Y. Shen,^{1,51} H. Y. Sheng,¹ X. Shi,^{1,47} X. D. Shi,^{59,47} J. J. Song,⁴⁰ Q. Q. Song,^{59,47} X. Y. Song,¹ Y. X. Song,^{37,‡} S. Sosio,^{62a,62c} C. Sowa,⁴ S. Spataro,^{62a,62c} F. F. Sui,⁴⁰ G. X. Sun,¹ J. F. Sun,¹⁶ L. Sun,⁶⁴ S. S. Sun,^{1,51} Y. J. Sun,^{59,47} Y. K. Sun,^{59,47} Y. Z. Sun,¹ Z. J. Sun,^{1,47} Z. T. Sun,¹ Y. X. Tan,^{59,47} C. J. Tang,⁴⁴ G. Y. Tang,¹ X. Tang,¹ V. Thoren,⁶³ B. Tsednee,²⁶ I. Uman,^{50d} B. Wang,¹ B. L. Wang,⁵¹ C. W. Wang,³⁵ D. Y. Wang,^{37,‡} K. Wang,^{1,47} L. L. Wang,¹ L. S. Wang,¹ M. Wang,⁴⁰ M. Z. Wang,^{37,‡} Meng Wang,^{1,51} P. L. Wang,¹ W. P. Wang,^{59,47} X. Wang,^{37,‡} X. F. Wang,³¹ X. L. Wang,^{9,‡‡} Y. Wang,⁴⁸ Y. Wang,^{59,47} Y. D. Wang,¹⁵ Y. F. Wang,^{1,47,51} Y. Q. Wang,¹ Z. Wang,^{1,47} Z. G. Wang,^{1,47} Z. Y. Wang,¹ Ziyi Wang,⁵¹ Zongyuan Wang,^{1,51} T. Weber,⁴ D. H. Wei,¹² P. Weidenkaff,²⁷ F. Weidner,⁵⁶ H. W. Wen,^{34,¶¶} S. P. Wen,¹ U. Wiedner,⁴ G. Wilkinson,⁵⁷ M. Wolke,⁶³ L. Wollenberg,⁴ L. H. Wu,¹ L. J. Wu,^{1,51} Z. Wu,^{1,47} L. Xia,^{59,47} S. Y. Xiao,¹ Y. J. Xiao,^{1,51} Z. J. Xiao,³⁴ Y. G. Xie,^{1,47} Y. H. Xie,⁶ T. Y. Xing,^{1,51} X. A. Xiong,^{1,51} G. F. Xu,¹ J. J. Xu,³⁵ Q. J. Xu,¹⁴ W. Xu,^{1,51} X. P. Xu,⁴⁵ F. Yan,⁶⁰ L. Yan,^{62a,62c} W. B. Yan,^{59,47} W. C. Yan,⁶⁷ W. C. Yan,² H. J. Yang,^{41,¶¶} H. X. Yang,¹ L. Yang,⁶⁴ R. X. Yang,^{59,47} S. L. Yang,^{1,51} Y. H. Yang,³⁵ Y. X. Yang,¹² Yifan Yang,^{1,51} Zhi Yang,²⁵ M. Ye,^{1,47} M. H. Ye,⁷ J. H. Yin,¹ Z. Y. You,⁴⁸ B. X. Yu,^{1,47,51} C. X. Yu,³⁶ J. S. Yu,²⁰ T. Yu,⁶⁰ C. Z. Yuan,^{1,51} X. Q. Yuan,^{37,‡} Y. Yuan,¹ C. X. Yue,³² A. Yuncu,^{50b,***} A. A. Zafar,⁶¹ Y. Zeng,²⁰ B. X. Zhang,¹ B. Y. Zhang,^{1,47} C. C. Zhang,¹ D. H. Zhang,¹ H. H. Zhang,⁴⁸ H. Y. Zhang,^{1,47} J. L. Zhang,⁶⁵ J. Q. Zhang,⁴ J. W. Zhang,^{1,47,51} J. Y. Zhang,¹ J. Z. Zhang,^{1,51} L. Zhang,¹ Lei Zhang,³⁵ S. F. Zhang,³⁵ T. J. Zhang,^{41,||} X. Y. Zhang,⁴⁰ Y. H. Zhang,^{1,47} Y. T. Zhang,^{59,47} Yan Zhang,^{59,47} Yao Zhang,¹ Yi Zhang,^{9,††} Yu Zhang,⁵¹ Z. H. Zhang,⁶ Z. P. Zhang,⁵⁹ Z. Y. Zhang,⁶⁴ G. Zhao,¹ J. Zhao,³² J. W. Zhao,^{1,47} J. Y. Zhao,^{1,51} J. Z. Zhao,^{1,47} Lei Zhao,^{59,47} Ling Zhao,¹

M. G. Zhao,³⁶ Q. Zhao,¹ S. J. Zhao,⁶⁷ T. C. Zhao,¹ Y. B. Zhao,^{1,47} Z. G. Zhao,^{59,47} A. Zhemchugov,^{28,‡} B. Zheng,⁶⁰ J. P. Zheng,^{1,47} Y. Zheng,^{37,†} Y. H. Zheng,⁵¹ B. Zhong,³⁴ L. Zhou,^{1,47} L. P. Zhou,^{1,51} Q. Zhou,^{1,51} X. Zhou,⁶⁴ X. K. Zhou,⁵¹ X. R. Zhou,^{59,47} A. N. Zhu,^{1,51} J. Zhu,³⁶ K. Zhu,¹ K. J. Zhu,^{1,47,51} S. H. Zhu,⁵⁸ W. J. Zhu,³⁶ X. L. Zhu,⁴⁹ Y. C. Zhu,^{59,47} Y. S. Zhu,^{1,51} Z. A. Zhu,^{1,51} J. Zhuang,^{1,47} B. S. Zou,¹ and J. H. Zou¹

(BESIII Collaboration)

- ¹*Institute of High Energy Physics, Beijing 100049, People's Republic of China*
²*Beihang University, Beijing 100191, People's Republic of China*
³*Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China*
⁴*Bochum Ruhr-University, D-44780 Bochum, Germany*
⁵*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*
⁶*Central China Normal University, Wuhan 430079, People's Republic of China*
⁷*China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China*
⁸*COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan*
⁹*Fudan University, Shanghai 200443, People's Republic of China*
¹⁰*G.I. Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*
¹¹*GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*
¹²*Guangxi Normal University, Guilin 541004, People's Republic of China*
¹³*Guangxi University, Nanning 530004, People's Republic of China*
¹⁴*Hangzhou Normal University, Hangzhou 310036, People's Republic of China*
¹⁵*Helmholtz Institute Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany*
¹⁶*Henan Normal University, Xinxiang 453007, People's Republic of China*
¹⁷*Henan University of Science and Technology, Luoyang 471003, People's Republic of China*
¹⁸*Huangshan College, Huangshan 245000, People's Republic of China*
¹⁹*Hunan Normal University, Changsha 410081, People's Republic of China*
²⁰*Hunan University, Changsha 410082, People's Republic of China*
²¹*Indian Institute of Technology Madras, Chennai 600036, India*
²²*Indiana University, Bloomington, Indiana 47405, USA*
^{23a}*INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*
^{23b}*INFN and University of Perugia, I-06100 Perugia, Italy*
^{24a}*INFN Sezione di Ferrara, I-44122 Ferrara, Italy*
^{24b}*University of Ferrara, I-44122 Ferrara, Italy*
²⁵*Institute of Modern Physics, Lanzhou 730000, People's Republic of China*
²⁶*Institute of Physics and Technology, Peace Ave. 54B, Ulaanbaatar 13330, Mongolia*
²⁷*Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany*
²⁸*Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia*
²⁹*Justus-Liebig-Universitaet Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany*
³⁰*KVI-CART, University of Groningen, NL-9747 AA Groningen, Netherlands*
³¹*Lanzhou University, Lanzhou 730000, People's Republic of China*
³²*Liaoning Normal University, Dalian 116029, People's Republic of China*
³³*Liaoning University, Shenyang 110036, People's Republic of China*
³⁴*Nanjing Normal University, Nanjing 210023, People's Republic of China*
³⁵*Nanjing University, Nanjing 210093, People's Republic of China*
³⁶*Nankai University, Tianjin 300071, People's Republic of China*
³⁷*Peking University, Beijing 100871, People's Republic of China*
³⁸*Qufu Normal University, Qufu 273165, People's Republic of China*
³⁹*Shandong Normal University, Jinan 250014, People's Republic of China*
⁴⁰*Shandong University, Jinan 250100, People's Republic of China*
⁴¹*Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China*
⁴²*Shanxi Normal University, Linfen 041004, People's Republic of China*
⁴³*Shanxi University, Taiyuan 030006, People's Republic of China*
⁴⁴*Sichuan University, Chengdu 610064, People's Republic of China*
⁴⁵*Soochow University, Suzhou 215006, People's Republic of China*
⁴⁶*Southeast University, Nanjing 211100, People's Republic of China*
⁴⁷*State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China*
⁴⁸*Sun Yat-Sen University, Guangzhou 510275, People's Republic of China*

- ⁴⁹*Tsinghua University, Beijing 100084, People's Republic of China*
^{50a}*Ankara University, 06100 Tandogan, Ankara, Turkey*
^{50b}*Istanbul Bilgi University, 34060 Eyup, Istanbul, Turkey*
^{50c}*Uludag University, 16059 Bursa, Turkey*
^{50d}*Near East University, Nicosia, North Cyprus, Mersin 10, Turkey*
⁵¹*University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China*
⁵²*University of Hawaii, Honolulu, Hawaii 96822, USA*
⁵³*University of Jinan, Jinan 250022, People's Republic of China*
⁵⁴*University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom*
⁵⁵*University of Minnesota, Minneapolis, Minnesota 55455, USA*
⁵⁶*University of Muenster, Wilhelm-Klemm-Str. 9, 48149 Muenster, Germany*
⁵⁷*University of Oxford, Keble Rd, Oxford OX13RH, United Kingdom*
⁵⁸*University of Science and Technology Liaoning, Anshan 114051, People's Republic of China*
⁵⁹*University of Science and Technology of China, Hefei 230026, People's Republic of China*
⁶⁰*University of South China, Hengyang 421001, People's Republic of China*
⁶¹*University of the Punjab, Lahore-54590, Pakistan*
^{62a}*University of Turin, I-10125 Turin, Italy*
^{62b}*University of Eastern Piedmont, I-15121 Alessandria, Italy*
^{62c}*INFN, I-10125, Turin, Italy*
⁶³*Uppsala University, Box 516, SE-75120 Uppsala, Sweden*
⁶⁴*Wuhan University, Wuhan 430072, People's Republic of China*
⁶⁵*Xinyang Normal University, Xinyang 464000, People's Republic of China*
⁶⁶*Zhejiang University, Hangzhou 310027, People's Republic of China*
⁶⁷*Zhengzhou University, Zhengzhou 450001, People's Republic of China*



(Received 7 August 2019; accepted 6 January 2020; published 5 February 2020; corrected 14 February 2020)

Using a sample of 1.31×10^9 J/ψ events collected with the BESIII detector, we perform a search for the rare decay $\eta' \rightarrow 4\pi^0$ via $J/\psi \rightarrow \gamma\eta'$. No significant η' signal is observed in the $4\pi^0$ invariant mass spectrum. With a Bayesian approach, the upper limit on the branching fraction is determined to be $\mathcal{B}(\eta' \rightarrow 4\pi^0) < 4.94 \times 10^{-5}$ at the 90% confidence level, which is a factor of 6 smaller than the previous experimental limit.

DOI: 10.1103/PhysRevD.101.032001

I. INTRODUCTION

The η' meson has a special role in improving the understanding of low-energy quantum chromodynamics

(QCD), and studies of its decays have attracted considerable theoretical [1] and experimental attention [2,3]. In addition to its important role in testing the fundamental

*Corresponding author.
changwl@ihep.ac.cn

†Also at the Novosibirsk State University, Novosibirsk, 630090, Russia.

‡Also at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People's Republic of China and School of Physics and Electronics, Hunan University, Changsha 410082, China.

§Also at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.

||Also at the Functional Electronics Laboratory, Tomsk State University, Tomsk, 634050, Russia.

¶Also at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People's Republic of China.

**Also at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany.

††Also at Istanbul Arel University, 34295 Istanbul, Turkey.

‡‡Also at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People's Republic of China.

§§Also at Harvard University, Department of Physics, Cambridge, Massachusetts, 02138, USA.

|||Also at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia.

¶¶Also at Ankara University, 06100 Tandogan, Ankara, Turkey.

***Also at Bogazici University, 34342 Istanbul, Turkey.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

discrete symmetries and searching for processes beyond the Standard Model (SM), η' decays offer unique opportunities to test the chiral perturbation theory (ChPT) [4] and the vector-meson dominance (VMD)[5] model.

In theory, $\eta' \rightarrow 4\pi^0$ is a highly suppressed decay because of the S-wave CP -violation. In the light of an effective chiral Lagrangian approach, the S-wave CP -violation in $\eta' \rightarrow 4\pi^0$ is induced by the so-called θ -term, which is an additional term in the QCD Lagrangian to account for the solution of the strong- CP problem. It was found that the S-wave CP -violation effect that contributed to this decay is at a level of 10^{-23} [6,7]. While higher-order contributions, involving a D-wave pion loop or the production of two f_2 tensor mesons (see Fig. 1), provide a CP -conserving route through which the decay can occur. By ignoring the tiny contribution from the latter process, calculations based on ChPT and VMD model predict the branching fraction caused by D-wave CP -conserving to be at the level of 10^{-8} [8]. However, the theoretical prediction is not strictly based on the effective field theory due to the lack of knowledge at such a high order in the chiral expansion and the use of a model to make an estimation. One does not know the reliability of that model *a priori*. Therefore, a search for the decay $\eta' \rightarrow 4\pi^0$ is useful to check the reliability of it.

So far the decay $\eta' \rightarrow 4\pi^0$ has not been observed. About three decades ago the first attempt to search for this mode was performed by the GAMS Collaboration, and the upper limit on the branching fraction was determined to be $\mathcal{B}(\eta' \rightarrow 4\pi^0) < 5 \times 10^{-4}$ at the 90% confidence level (C.L.) [9]. The more recent upper limit of 3.2×10^{-4} at 90% C.L. was obtained by the GAMS- 4π Collaboration [10].

Although η' meson can not be produced directly in e^+e^- annihilations, the decay $J/\psi \rightarrow \gamma\eta'$, with a branching




FIG. 1. D-wave pion-loop (a) and intermediate f_2 mesons contribution (b) to $\eta' \rightarrow 4\pi^0$ [8].

fraction of $(5.13 \pm 0.17) \times 10^{-3}$ [11], provides an abundant source of η' meson in this environment. The BESIII experiment has exploited this production mode to perform a series of studies of η' decays [12], based on a sample of $(1310.6 \pm 7.0) \times 10^6$ J/ψ events taken at the center-of-mass energy of 3.097 GeV [13] with the BESIII detector, corresponding to 6.7×10^6 η' events. In this paper, using this same J/ψ sample, we perform a search for $\eta' \rightarrow 4\pi^0$ via $J/\psi \rightarrow \gamma\eta'$.

II. BESIII DETECTOR AND MONTE CARLO SIMULATION

The BESIII detector [14] is a magnetic spectrometer located at the Beijing Electron Positron Collider (BEPCII) [15], which is a double-ring e^+e^- collider with a design peak luminosity of $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ at the center-of-mass energy of 3.773 GeV. The cylindrical core of the BESIII detector consists of a helium-based multilayer drift chamber (MDC), a plastic scintillator time-of-flight system (TOF), and a CsI(Tl) electromagnetic calorimeter (EMC), which are all enclosed in a superconducting solenoidal magnet providing a 1.0 T (0.9 T in 2012) magnetic field. The solenoid is supported by an octagonal flux-return yoke with resistive plate counter muon-identifier modules interleaved with steel. The acceptance of charged particles and photons is 93% over the 4π solid angle. The charged-particle momentum resolution at 1 GeV/c is 0.5%, and the dE/dx resolution is 6% for electrons from Bhabha scattering. The EMC measures photon energies with a resolution of 2.5% (5%) at 1 GeV in the barrel (end cap) region. The time resolution of the TOF barrel part is 68 ps, while that of the end cap part is 110 ps.

Simulated samples produced with a GEANT4-based [16] Monte Carlo (MC) simulation framework, which includes the geometric description of the BESIII detector and the detector response, are used to determine the detection efficiency and to estimate the backgrounds. The simulation includes the beam energy spread and initial state radiation (ISR) in e^+e^- annihilation modeled with the generator KKMC [17]. The inclusive MC sample consists of the production of the J/ψ resonance, and the continuum processes incorporated in KKMC [17]. The known decay modes are modeled with [18,19] using branching fractions taken from the Particle Data Group (PDG) [20], and the remaining unknown decays of the charmonium states with LUNDCHARM [21]. The final state radiation (FSR) from charged final-state particles are incorporated with the PHOTOS package [22].

III. EVENT SELECTION

In this analysis, the pseudoscalar mesons η' and π^0 are reconstructed in the modes $\eta' \rightarrow 4\pi^0$ and $\pi^0 \rightarrow \gamma\gamma$. Candidate $J/\psi \rightarrow \gamma 4\pi^0$ decays are chosen by selecting




FIG. 2. χ^2_{8C} distribution of candidate events in the η' signal region showing data and the contribution from the considered background sources. The arrow indicates the selection requirement of $\chi^2_{8C} < 30$.

events with at least nine isolated photons and no charged track. Photon candidates are reconstructed from clusters of energy deposited in the EMC. Photon candidates are required to have at least 25 MeV of energy for barrel showers ($|\cos\theta| \leq 0.8$) or 50 MeV for end cap showers ($0.86 \leq |\cos\theta| \leq 0.92$).

The photon candidate with the maximum energy deposited in the EMC is treated as the radiative photon directly originating from the J/ψ decay. For the two-body signal decay $J/\psi \rightarrow \gamma\eta'$, this photon carries a unique energy of 1.4 GeV. To reconstruct the π^0 candidate from the remaining photons, a one-constraint (1C) kinematic fit is performed to each photon pair with the invariant mass constrained to the π^0 mass, with the requirement that the goodness-of-fit $\chi^2_{1C}(\gamma\gamma) < 25$. And at least four π^0 candidates are required for further analysis. Then an eight-constraint (8C) kinematic fit is performed for the $\gamma\pi^0\pi^0\pi^0$ combination by enforcing energy-momentum conservation and constraining the invariant masses of each of the four photon pairs to the nominal π^0 mass. If more than one combination is found in an event, only the one with the smallest χ^2_{8C} is retained. The χ^2_{8C} distribution is shown in Fig. 2. Candidate events with $\chi^2_{8C} > 30$ are rejected. The MC study shows that after the above selection, about 8% of events have a miscombination of photons, which mainly occurs between the different π^0 candidates. Since this is not the miscombination between the radiative photon and the photon from π^0 candidates, the invariant mass of $4\pi^0$ is still in the η' mass region.

To ensure a good description of data, a signal MC simulation is modeled with the decay amplitudes in Ref. [8], which assumes the pion-loop contribution as shown in Fig. 1(a) [the contribution from f_2 mesons shown in Fig. 1(b) is considered to be much smaller]. After the full event selection, the detection efficiency (ϵ) is determined to be $(2.46 \pm 0.01)\%$.

Figure 3 shows the $4\pi^0$ mass spectrum, $M(4\pi^0)$, after selection. No significant η' signal is evident. In order to




FIG. 3. The $M(4\pi^0)$ distribution in data, together with the total fit result and the contributions from nonpeaking background and the peaking background $J/\psi \rightarrow \gamma\eta'$, $\eta' \rightarrow \pi^0\pi^0\eta$, $\eta \rightarrow \pi^0\pi^0\pi^0$. Also shown is the expected shape of the signal contribution, with arbitrary normalization.

investigate possible sources of contamination, we apply the selection to an inclusive MC sample of 1.22×10^9 J/ψ events. Since the decay $J/\psi \rightarrow n\pi^0$, where n is the number of π^0 mesons, is forbidden because of charge conjugation conservation, we find that the background comes mainly from decays with the same final state as the signal, for example, $J/\psi \rightarrow \eta\omega$, $\eta \rightarrow \pi^0\pi^0\pi^0$, $\omega \rightarrow \gamma\pi^0$ or from radiative decays with more than four π^0 's in the final states, of which the dominant mode is $J/\psi \rightarrow \gamma\eta'$, $\eta' \rightarrow \pi^0\pi^0\eta$, $\eta \rightarrow \pi^0\pi^0\pi^0$. None of these background events peaks in the η' mass region, while the background events from $J/\psi \rightarrow \gamma\eta'$ ($\eta' \rightarrow \pi^0\pi^0\eta$) contributes to a broad structure around 0.88 GeV/ c^2 in the $4\pi^0$ mass spectrum. Then the dedicated exclusive MC samples are generated for these background channels. The decays of $J/\psi \rightarrow \gamma\eta'$ and $J/\psi \rightarrow \eta\omega$, are generated using a helicity amplitude model [19], where the cascade decays of $\eta' \rightarrow \pi^0\pi^0\eta$ and $\eta' \rightarrow 3\pi^0$ are modeled with the Dalitz plot analysis results presented in Refs. [23,24]. And the uniform phase space events are generated for the other background contributions. In accordance with the branching fractions and the

TABLE I. The main background channels and their expected contribution to the selected sample.

| Decay mode | Generated events | Normalized events |
|--|-------------------|-------------------|
| $J/\psi \rightarrow \gamma\eta', \eta' \rightarrow \pi^0\pi^0\eta, \eta \rightarrow \pi^0\pi^0\pi^0$ | 1.5×10^7 | 496 ± 18 |
| $J/\psi \rightarrow \eta\omega, \eta \rightarrow \pi^0\pi^0\pi^0, \omega \rightarrow \gamma\pi^0$ | 1×10^7 | 131 ± 4 |
| $J/\psi \rightarrow \gamma\pi^0\pi^0\eta, \eta \rightarrow \pi^0\pi^0\pi^0$ | 2.3×10^7 | 38 ± 2 |
| $J/\psi \rightarrow \gamma\pi^0\eta, \eta \rightarrow \pi^0\pi^0\pi^0$ | 3×10^6 | 24 ± 1 |
| $J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi^0 a^0$ | 1.2×10^7 | 11 ± 1 |
| $J/\psi \rightarrow \gamma f_2(1270), f_2(1270) \rightarrow \pi^0\pi^0\pi^0$ | 2×10^6 | 10 ± 1 |
| $J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi^0\pi^0\eta$ | 1.2×10^7 | 5 ± 1 |

reconstruction efficiencies, the normalized background events are listed in Table I, and the expected peaking contribution of $J/\psi \rightarrow \gamma\eta'$ ($\eta' \rightarrow \pi^0\pi^0\eta$) to the $4\pi^0$ mass spectrum is displayed in Fig. 3.

IV. FIT MODEL AND UPPER LIMIT ON BRANCHING FRACTION OF $\eta' \rightarrow 4\pi^0$

An unbinned maximum likelihood fit is performed on the $4\pi^0$ mass spectrum, allowing for background contributions and a possible signal component. In the fit, the line shape of the η' signal is determined by a MC simulation. The shape of peaking background $J/\psi \rightarrow \gamma\eta', \eta' \rightarrow \pi^0\pi^0\eta, \eta \rightarrow \pi^0\pi^0\pi^0$ is obtained from the dedicated MC simulation [23,24] with the number fixed to the normalized value while the nonpeaking background contribution is described by a third-order Chebychev polynomial function with the number and parameters of Chebychev function free.

A Bayesian approach is used to determine an upper limit on the branching fraction of $\eta' \rightarrow 4\pi^0$. A series of unbinned extended maximum likelihood fits are performed for different assumed values of the signal yield N , and for each fit the negative log-likelihood \mathcal{S} is determined. For each value of N the branching fraction is

$$\mathcal{B}(\eta' \rightarrow 4\pi^0) = \frac{N}{N_{J/\psi} \cdot \varepsilon \cdot \mathcal{B}(J/\psi \rightarrow \gamma\eta') \cdot \mathcal{B}(\pi^0 \rightarrow \gamma\gamma)^4}, \quad (1)$$

where $N_{J/\psi} = (1310.6 \pm 7.0) \times 10^6$ is the number of J/ψ events [13], ε is the detection efficiency, $\mathcal{B}(J/\psi \rightarrow \gamma\eta')$ and $\mathcal{B}(\pi^0 \rightarrow \gamma\gamma)$ are the branching fractions of $J/\psi \rightarrow \gamma\eta'$ and $\pi^0 \rightarrow \gamma\gamma$, respectively, which are taken from Ref. [11].

The distribution of normalized likelihood values, defined as $\mathcal{L}(B) = \exp(-[\mathcal{S}(B) - \mathcal{S}_{\min}])$, where \mathcal{S}_{\min} is the lowest negative log-likelihood obtained from the ensemble of fits, is taken as the probability density function (PDF) for the expected branching fraction of $\eta' \rightarrow 4\pi^0$. The upper limit on the branching fraction at the 90% C.L., defined as \mathcal{B}_{UL} , corresponds to the branching fraction at 90% of the integral of the PDF,

$$\frac{\int_0^{\mathcal{B}_{UL}} \mathcal{L}(B) dB}{\int_0^\infty \mathcal{L}(B) dB} = 0.9, \quad (2)$$

and is found to be 4.57×10^{-5} , considering statistical uncertainties alone.

V. SYSTEMATIC UNCERTAINTIES

Two categories of systematic uncertainty are considered: those associated with the fit model and procedure, and those which enter when using Eq. (1) to express the signal yield as a branching fraction.

The fit-related uncertainties come mainly from the fitting ranges, signal shape, nonpeaking background shape,

peaking background shape and the number of the peaking background events.

The uncertainties due to the fit range are considered by varying the fit ranges of 5 MeV/c² or 10 MeV/c². And the maximum change on the results is taken as the systematic uncertainty from fit range. In the fit to the $M(4\pi^0)$ distribution, signal shape is taken from the MC simulation. To assess the uncertainty due to the signal shape, an alternative fit is performed by convolving a Gaussian function with a fixed resolution of 2.6 MeV and mean of 2.1 MeV which are obtained from a high purity control sample of $J/\psi \rightarrow \gamma\eta', \eta' \rightarrow \pi^0\pi^0\eta, \eta \rightarrow \gamma\gamma$. The uncertainty from the nonpeaking background shape is determined by using a fourth-order Chebychev polynomial in place of the third-order Chebychev polynomial. To assess the uncertainty associated with the number of the peaking background events, its contribution is recalculated after varying the branching fractions of $J/\psi \rightarrow \gamma\eta'$ and its cascade decays, $\eta' \rightarrow \pi^0\pi^0\eta$ and $\eta \rightarrow \pi^0\pi^0\pi^0$, within their uncertainties, and new fits are performed. The systematic uncertainty associated with peaking background shape is evaluated by convolving a Gaussian function with the resolution and mean value left free. Among these cases, the dominant fit-model uncertainty arises from fitting range [0.705, 1.095] GeV/c², and it changes the upper limit at the 90% C.L. to $\mathcal{B}_{UL} = 4.88 \times 10^{-5}$.

The other category of systematic uncertainties, summarized in Table II, has contributions from the knowledge of the photon detection efficiency, the efficiency of the kinematic fit, signal model, MC statistical uncertainty for the detection efficiency, the branching fractions of the subdecays involved in the signal process, and the total number of J/ψ events.

The uncertainty from the photon detection is investigated with a high purity control sample of $J/\psi \rightarrow \pi^+\pi^-\pi^0$. It is found that the differences between data and MC simulation are 0.5% and 1.5% for each photon deposited in the barrel and end cap of the EMC, respectively. With the same approach as used in Ref. [25], the uncertainty on the detection efficiency for each photon in the signal decay is estimated to be 0.53%, and thus the nine photons in the final state induce an overall uncertainty of 4.8%.

TABLE II. Summary of the systematic uncertainties unrelated to the fit model. For each component the relative impact on the branching fraction is listed in %.

| Source | Systematic uncertainties |
|---|--------------------------|
| Photon detection | 4.8 |
| Kinematic fit | 4.1 |
| Signal model | 2.0 |
| $\mathcal{B}(J/\psi \rightarrow \gamma\eta')$ | 3.1 |
| $\mathcal{B}(\pi^0 \rightarrow \gamma\gamma)$ | 0.12 |
| MC statistic | 0.4 |
| Number of J/ψ events | 0.54 |
| Total | 7.3 |

The uncertainty associated with the kinematic fit is estimated by adjusting the components of the photon-energy error matrix in the signal MC sample to reflect the known difference in the resolution between data and the MC simulation [26]. From the study of the $\psi(3686) \rightarrow \gamma\chi_{c1}(\chi_{c1} \rightarrow 4\pi^0)$ decay [27], it is known that the energy resolution in data is 4% wider than in a MC simulation. The relative difference in efficiency, 4.1%, is taken as the systematic uncertainty from the kinematic fit.

In the normal fit, the cascade decay $\eta' \rightarrow 4\pi^0$ is described with the decay amplitude in Ref. [8]. A fit with an alternative signal model replacing $\eta' \rightarrow 4\pi^0$ decay amplitude with a phase space (PHSP) distribution is performed. The change of the efficiency, 2.0%, is taken as the uncertainty due to the signal model.

The relative uncertainty in the knowledge of the branching fractions of $J/\psi \rightarrow \gamma\eta'$ and $\pi^0 \rightarrow \gamma\gamma$ [11] induces a corresponding uncertainty on the calculated upper limit on the branching fraction. The uncertainty of the detection efficiency, 0.4%, caused by MC statistics is also taken as a source of systematic uncertainty. The number of J/ψ events is determined from the measured number of hadronic decays and is found to be $(1310.6 \pm 7.0) \times 10^6$ [13], which corresponds to a relative uncertainty of 0.54%.

Assuming all systematic uncertainties presented in Table II are independent, the total relative uncertainty is obtained to be 7.3%, by adding all individual uncertainties in quadrature.

VI. RESULT

The final upper limit on the branching fraction is determined by convolving the likelihood distribution \mathcal{L} with the systematic uncertainties to obtain the smeared likelihood $\mathcal{L}^{\text{smeared}}$,

$$\mathcal{L}^{\text{smeared}}(\mathcal{B}) = \int \mathcal{L}\left(\frac{\varepsilon}{\bar{\varepsilon}}\mathcal{B}\right) \exp\left(-\frac{(\varepsilon - \bar{\varepsilon})^2}{2\sigma_{\varepsilon}^2}\right) d\varepsilon. \quad (3)$$

In this exercise all components listed in Table II, whatever their nature, can be considered as an uncertainty on the




FIG. 4. Normalized likelihood distribution before (black dots) and after (red stars) convolution with systematic uncertainty.

detection efficiency ε . The nominal efficiency value is $\bar{\varepsilon}$, σ_{ε} is the absolute total systematic uncertainty on the efficiency, and \mathcal{B} is the branching fraction of $\eta' \rightarrow 4\pi^0$.

Figure 4 shows the normalized likelihood distribution after taking all systematic uncertainties into account. The corresponding upper limit of the branching fraction of $\eta' \rightarrow 4\pi^0$ at the 90% C.L. is determined to be 4.94×10^{-5} .

VII. SUMMARY

Using a sample of 1.31×10^9 J/ψ events collected with the BESIII detector, a search for the decay $\eta' \rightarrow 4\pi^0$ is performed via $J/\psi \rightarrow \gamma\eta'$. No evidence for the rare decay $\eta' \rightarrow 4\pi^0$ is found, and an upper limit of $\mathcal{B}(\eta' \rightarrow 4\pi^0) < 4.94 \times 10^{-5}$ is set at the 90% confidence level. This limit is approximately a factor of 6 smaller than the previous most stringent result [10].

The current limit is still far to reach the theoretical predication with a level of 10^{-8} [8]. Further studies of η' rare decays are still necessary to test the ChPT and VMD model and look for the CP -violation (S-wave) $\eta' \rightarrow 4\pi^0$ decay. A sample of 10^{10} J/ψ events has now been collected at BESIII, which will allow for even more sensitive searches to be performed for this important decay mode.

ACKNOWLEDGMENTS

The BESIII Collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key Basic Research Program of China under Contract No. 2015CB856700; National Natural Science Foundation of China (NSFC) under Contracts No. 11335008, No. 11425524, No. 11625523, No. 11635010, No. 11675184, No. 11735014; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contracts No. U1532257, No. U1532258, No. U1732263, No. U1832207; CAS Key Research Program of Frontier Sciences under Contracts No. QYZDJ-SSW-SLH003, No. QYZDJ-SSW-SLH040; 100 Talents Program of CAS; the Institute of Nuclear and Particle Physics and Shanghai Key Laboratory for Particle Physics and Cosmology; ERC under Contract No. 758462; German Research Foundation DFG under Contracts No. Collaborative Research Center CRC 1044, No. FOR 2359; Istituto Nazionale di Fisica Nucleare, Italy; Koninklijke Nederlandse Akademie van Wetenschappen (KNAW) under Contract No. 530-4CDP03; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Science and Technology fund; STFC (United Kingdom); The Knut and Alice Wallenberg Foundation (Sweden) under Contract No. 2016.0157; The Royal Society, UK under Contracts No. DH140054, No. DH160214; The Swedish Research Council; U.S. Department of Energy under Contracts No. DE-FG02-05ER41374, No. DE-SC-0010118, No. DE-SC-0012069; University of Groningen (RuG) and the Helmholtzzentrum fuer Schwerionenforschung GmbH (GSI), Darmstadt.

- [1] H. Nagahiro, M. Takizawa, and S. Hirenzaki, *Phys. Rev. C* **74**, 045203 (2006).
- [2] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **97**, 012003 (2018).
- [3] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **112**, 251801 (2014).
- [4] S. Scherer, *Adv. Nucl. Phys.* **27**, 277 (2003).
- [5] J. J. Sakurai, *Currents and Mesons* (University of Chicago Press, Chicago, 1969).
- [6] A. Pich and E. de Rafael, *Nucl. Phys.* **B367**, 313 (1991).
- [7] K. Ottnad, B. Kubis, and U. G. Meißner, and F. K. Guo, *Phys. Lett. B* **687**, 42 (2010).
- [8] F. K. Guo, B. Kubis, and A. Wirzba, *Phys. Rev. D* **85**, 014014 (2012).
- [9] D. Alde *et al.*, *Z. Phys. C* **36**, 603 (1987).
- [10] S. V. Donskov, V. N. Kolosov, A. A. Lednev, Y. V. Mikhailov, V. A. Polyakov, V. D. Samoylenko, and G. V. Khaustov, *Mod. Phys. Lett. A* **29**, 1450213 (2014).
- [11] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018).
- [12] S. S. Fang, A. Kupsc, and D. H. Wei, *Chin. Phys. C* **42**, 042002 (2018).
- [13] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **41**, 013001 (2017).
- [14] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 345 (2010).
- [15] C. H. Yu *et al.*, *Proceedings of IPAC2016, Busan, Korea* (2016), <https://dx.doi.org/10.18429/JACoW-IPAC2016-TUYA01>.
- [16] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [17] S. Jadach, B. F. L. Ward, and Z. Was, *Phys. Rev. D* **63**, 113009 (2001); *Comput. Phys. Commun.* **130**, 260 (2000).
- [18] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [19] R. G. Ping, *Chin. Phys. C* **32**, 599 (2008).
- [20] C. Patrignani *et al.* (Particle Data Group), *Chin. Phys. C* **40**, 100001 (2016).
- [21] J. C. Chen, G. S. Huang, X. R. Qi, D. H. Zhang, and Y. S. Zhu, *Phys. Rev. D* **62**, 034003 (2000); R. L. Yang, R. G. Ping, and H. Chen, *Chin. Phys. Lett.* **31**, 061301 (2014).
- [22] E. Richter-Was, *Phys. Lett. B* **303**, 163 (1993).
- [23] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **97**, 012003 (2018).
- [24] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **92**, 012014 (2015).
- [25] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **96**, 112012 (2017).
- [26] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **94**, 072005 (2016).
- [27] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **83**, 012006 (2011).

Correction: Contract numbers were missing in the Acknowledgments section and have been inserted.